


AMENDMENTS TO THE CLAIMS

1. (Previously Presented) A pyrimidine of the formula I



in which the index and the substituents are as defined below:

- n is an integer from 1 to 5;
- L is halogen, cyano, nitro, cyanato (OCN), C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=S)-N(A')A, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A;
- m is 0, 1 or 2;
- A, A', A'' independently of one another are hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl, where the organic radicals may be partially or fully halogenated or may be substituted by cyano or C₁-C₄-alkoxy, or A and A' together with the atoms to which they are attached are a five- or six-

membered saturated, partially unsaturated or aromatic heterocycle
which contains one to four heteroatoms from the group consisting
of O, N and S;

R¹ is C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₃-C₁₂-cycloalkyl, C₃-C₁₀-
cycloalkenyl;

R² is halogen, cyano, C₁-C₄-alkoxy, C₃-C₄-alkenyloxy or C₃-C₄-alkynyloxy;

R³ is a five- or six-membered saturated, partially unsaturated or aromatic mono- or
bicyclic heterocycle which contains one to four heteroatoms from the group
consisting of O, N and S,

where the aliphatic, alicyclic or aromatic groups of the radical definitions of L, R¹, R²
and/or R³ for their part may be partially or fully halogenated or may carry one to four
groups R^a:

R^a is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy,
C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, OH, SH, two vicinal groups R^a may be
(=O) or (=S), C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-
cycloalkenyloxy, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA),
N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or

$S(=O)_m-N(A')A$, where m , A , A' , A'' are as defined above and where the aliphatic, alicyclic or aromatic groups for their part may be partially or fully halogenated or may carry one to three groups R^b , where R^b has the same meaning as R^a .

2. (Previously Presented) A pyrimidine as claimed in claim 1, in which the index and the substituents are as defined below:

L is halogen, cyano, C_1 - C_8 -alkyl, C_2 - C_{10} -alkenyl, C_2 - C_{10} -alkynyl, C_1 - C_6 -alkoxy, C_2 - C_{10} -alkenyloxy, C_2 - C_{10} -alkynyloxy, $-C(=O)-O-A$, $N(A')-C(=O)-A$ or $S(=O)_m-A$;

m is 0, 1 or 2;

A, A', A'' independently of one another are hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, where the organic radicals may be partially or fully halogenated or A and A' together with the atoms to which they are attached are a partially unsaturated or aromatic heterocycle which contains one to four heteroatoms from the group consisting of O, N and S;

R^1 is C_1 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_2 - C_{10} -alkynyl, C_3 - C_{12} -cycloalkyl,

C₃-C₁₀-cycloalkenyl;

R² is cyano or chlorine,

where the aliphatic, alicyclic or aromatic groups of the radical definitions of L, R¹ and/or R³ for their part may be partially or fully halogenated or may carry one to four groups R^a:


R^a is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A.

3. (Original) A pyrimidine as claimed in claim 1, in which R³ is pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, oxazolyl, isoxazolyl, 1,3,4-oxadiazolyl, furanyl, thiophenyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, pyrrolidinyl, piperidinyl, hexahydroazepinyl or dihydropyridinyl, where the heterocycle may be attached to the pyrimidine ring via carbon or nitrogen and may carry up to three substituents R^a:

R^a is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, OH, SH, two vicinal groups R^a may be (=O)

or (=S), C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A.

4. (Original) A pyrimidine as claimed in claim 1, in which R³ is pyrazol-1-yl, [1,2,4]-triazol-1-yl, pyridin-2-yl, pyrimidin-2-yl, pyridazin-3-yl, pyrrolidin-2-on-1-yl, piperidin-2-on-1-yl, hexahydro-2H-azepin-2-on-1-yl, pyrrolidin-2-thion-1-yl, piperidin-2-thion-1-yl, hexahydro-2H-azepin-2-thion-1-yl, 1,2-dihydropyridin-2-on-1-yl.
5. (Previously Presented) A pyrimidine as claimed in claim 1, in which R² is chlorine.
6. (Previously Presented) A pyrimidine as claimed in any of claims 1 to 5, in which the phenyl group substituted by L_n is the group B




where # is the point of attachment to the pyrimidine skeleton and
L¹ is fluorine, chlorine, CH₃ or CF₃;

L^2, L^4 independently of one another are hydrogen, CH_3 or fluorine;

L^3 is hydrogen, fluorine, chlorine, bromine, cyano, CH_3 , SCH_3 , OCH_3 , SO_2CH_3 , $CO-NH_2$, $CO-NHCH_3$, $CO-NHC_2H_5$, $CO-N(CH_3)_2$, $NH-C(=O)CH_3$, $N(CH_3)-C(=O)CH_3$ or $COOCH_3$ and


L^5 is hydrogen, fluorine, chlorine or CH_3 .

7. (Previously Presented) A process for preparing a pyrimidine of the formula I as claimed in claim 1, where R^3 is a nitrogen-containing heterocycle attached via nitrogen, which comprises reacting a compound of the formula III,



in which the substituents L_n , R^1 and R^2 are as defined in claim 1 and X is halogen, C_1 - C_6 -alkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfoxyl or C_1 - C_6 -alkylsulfenyl, with a heterocycle of the formula R^3-H (IV), optionally in the presence of a base.

8. (Currently Amended) An intermediate of the formula III



wherein:

R¹ is C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₃-C₁₂-cycloalkyl, C₃-C₁₀-cycloalkenyl;

L_n is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, -C(=O)-O-A, N(A')-C(=O)-A or S(=O)_m-


A;

m is 0, 1 or 2;

X is halogen, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylsulfoxyl or C₁-C₆-alkylsulfenyl, with a heterocycle of the formula R³-H (IV), optionally in the presence of a base,

~~in which the substituent R¹ is as defined in claim 1, L_n is as defined in claim 2, X is as defined in~~
~~claim 7~~ and R² is cyano, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-alkoxy, C₃-C₄-alkenyloxy or C₃-C₄-alkynyloxy, where the alkyl, alkenyl and alkynyl radicals of R² may be substituted by halogen, cyano, nitro, C₁-C₂-alkoxy or C₁-C₄-alkoxycarbonyl.

9. (Currently Amended) A pesticidal composition, which comprises a solid or liquid carrier and ~~a compound of the formula I as claimed in claim 1~~ a pyrimidine of the formula I



in which the index and the substituents are as defined below:

n is an integer from 1 to 5;

L is halogen, cyano, nitro, cyanato (OCN), C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=S)-N(A')A, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A;

m is 0, 1 or 2;

A, A', A'' independently of one another are hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl, where the organic radicals may be partially or fully halogenated or

may be substituted by cyano or C₁-C₄-alkoxy, or A and A' together with the atoms to which they are attached are a five- or six-membered saturated, partially unsaturated or aromatic heterocycle which contains one to four heteroatoms from the group consisting of O, N and S;

R¹ is C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₃-C₁₂-cycloalkyl, C₃-C₁₀-cycloalkenyl;

R² is halogen, cyano, C₁-C₄-alkoxy, C₃-C₄-alkenyloxy or C₃-C₄-alkynyloxy;


R³ is a five- or six-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocycle which contains one to four heteroatoms from the group consisting of O, N and S,

where the aliphatic, alicyclic or aromatic groups of the radical definitions of L, R¹, R² and/or R³ for their part may be partially or fully halogenated or may carry one to four groups R^a;

R^a is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, OH, SH, two vicinal groups R^a may be (=O) or (=S), C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -

C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A, where m, A, A', A'' are as defined above and where the aliphatic, alicyclic or aromatic groups for their part may be partially or fully halogenated or may carry one to three groups R^b, where R^b has the same meaning as R^a.

10. (Currently Amended) A method for controlling phytopathogenic harmful fungi, which comprises treating the fungi or the materials, plants, the soil or seeds to be protected against fungal attack with an effective amount of ~~a compound of the formula I as claimed in claim 1~~ a pyrimidine of the formula I



in which the index and the substituents are as defined below:

n is an integer from 1 to 5;

L is halogen, cyano, nitro, cyanato (OCN), C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=S)-N(A')A,

-C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A,
N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or
S(=O)_m-N(A')A;

m is 0, 1 or 2;

A, A', A'' independently of one another are hydrogen, C₁-C₆-alkyl, C₂-C₆-
alkenyl, C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl,
where the organic radicals may be partially or fully halogenated or
may be substituted by cyano or C₁-C₄-alkoxy, or A and A' together
with the atoms to which they are attached are a five- or six-
membered saturated, partially unsaturated or aromatic heterocycle
which contains one to four heteroatoms from the group consisting
of O, N and S;

R¹ is C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₃-C₁₂-cycloalkyl, C₃-C₁₀-
cycloalkenyl;

R² is halogen, cyano, C₁-C₄-alkoxy, C₃-C₄-alkenyloxy or C₃-C₄-alkynyloxy;

R³ is a five- or six-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocycle which contains one to four heteroatoms from the group consisting of O, N and S,

where the aliphatic, alicyclic or aromatic groups of the radical definitions of L, R¹, R² and/or R³ for their part may be partially or fully halogenated or may carry one to four groups R^a:

R^a is halogen, cyano, C₁-C₈-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₁-C₆-alkoxy, C₂-C₁₀-alkenyloxy, C₂-C₁₀-alkynyloxy, OH, SH, two vicinal groups R^a may be (=O) or (=S), C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₃-C₆-cycloalkoxy, C₃-C₆-cycloalkenyloxy, -C(=O)-A, -C(=O)-O-A, -C(=O)-N(A')A, C(A')(=N-OA), N(A')A, N(A')-C(=O)-A, N(A'')-C(=O)-N(A')A, S(=O)_m-A, S(=O)_m-O-A or S(=O)_m-N(A')A, where m, A, A', A'' are as defined above and where the aliphatic, alicyclic or aromatic groups for their part may be partially or fully halogenated or may carry one to three groups R^b, where R^b has the same meaning as R^a.

11. (Previously Presented) A pyrimidine as claimed in claim 1, wherein R² is halogen, cyano or C₁-C₄-alkoxy.
12. (Previously Presented) A pyrimidine as claimed in claim 1, wherein R¹ is C₃-C₈-alkyl, C₃-C₈-alkenyl, C₃-C₈-alkynyl, C₃-C₆-cycloalkyl or C₅-C₆-cycloalkenyl.

13. (Currently Amended) A pyrimidine as claimed in claim 1, wherein R^1 is C_1 - C_6 -alkyl or ~~C_4 - C_6 -haloalkyl~~.

14. (Previously Presented) A pyrimidine as claimed in claim 1, wherein R^1 is selected from the group consisting of 2-methyl-butyl, cyclohexyl, but-1-en-4-yl, methyl, 3-methyl-but-1-enyl, 2-hydroxy-3-methyl-butyl, and 2-methyl-propyl.

15. (Previously Presented) The pyrimidine of claim 14, wherein R^2 is halogen, cyano or C_1 - C_4 -alkoxy.

16. (Previously Presented) The pyrimidine of claim 15, wherein R^3 is selected from the group consisting of [1,2,4] triazol-1-yl, pyrazol-1-yl, 1,2,3-triazol-1-yl, 3-cyano-1,2,4-triazol-1-yl, 7-amino-indazol-1-yl, and 3-amino-pyrazol-1-yl.